



Sulfoxaflor and bees

Sulfoxaflor has recently come onto the market in China (Shao et al. 2013) and the USA (Dow Agro Sciences 2013; USEPA 2013), was reviewed by the EFSA in 2014 (EFSA 2014) and approved in the European Union in 2015.

Sulfoxaflor, an active substance of the family of sulfoximines, is a fourth-generation neonicotinoid based on its target receptors and binding sites, that exhibits a high insecticidal activity against a broad range of sap-feeding insects (Babcock et al. 2011, Cutler et al. 2013).

Mode of action

It acts on nAChRs, with a pharmacological profile similar to that of imidacloprid (Cutler et al. 2013). For this reason, sulfoxaflor may be considered as a neonicotinoid. However, the nature of the interactions with nAChRs differs between sulfoxaflor and the other neonicotinoids (Sparks et al. 2013). The action of sulfoxaflor and other sulfoximines, similar to that of imidacloprid, involves receptor desensitization, receptor selectivity, a differential action at low and high doses and, probably, receptor desensitization after a prolonged exposure (Oliveira et al. 2011, Watson et al. 2011, Cutler et al. 2013). This needs to be taken into account when considering possibilities for insecticide rotation in order to manage resistance toward neonicotinoids based on modification of the target site (Cutler et al. 2013).

Considering the pharmacology (binding sites at nACh receptors) of sulfoxaflor, insects belonging to the hemiptera order show the highest sensitivity. However, toxicity to non-hemiptera insects can be also expected at higher application rates (Cutler et al. 2013).

Metabolism

Compared to imidacloprid, acetamiprid, dinotefuran, thiamethoxam, and clothianidin for which the extents of metabolism are respectively 85.1, 95.5, 55.1, 46.8, and 45.6 % after 24 h, sulfoxaflor presents an almost undetectable metabolism (Sparks et al. 2012).

Toxicity to bees

Table in Annex 1 summarises the LD₅₀ values included in the European dossier. The EFSA identified a high risk to bees and several data gaps were also pointed out, namely the effects on non-*Apis* bees or the risk of field uses when applying risk mitigation measures. Among others, potential adverse effects on bee brood could not be excluded together with toxicity following foliar application (EFSA 2014).

The USEPA determines that the acute oral toxicity (LD₅₀) to be 0.052 µg of sulfoxaflor per bee and that the acute contact toxicity (LD₅₀) to be 0.13 µg of active substance per bee. In parallel, the EFSA determines an oral LD₅₀ of 0.146 µg of sulfoxaflor per bee and a contact LD₅₀ of 0.379 µg of sulfoxaflor per bee. When sulfoxaflor is included into commercial formulations the oral LD₅₀ of

two products is 0.065 or 0.075 µg of sulfoxaflor per bee, respectively. This may indicate possible synergistic effects with some of the coformulants. Sulfoxaflor has proved to potentiate the herbicidal effects of pyridine carboxylic acid compounds (Satchivi and Schmitzer 2011), however it is not clear to what extend the insecticidal effect of sulfoxaflor is also enhanced by this combination.

The EFSA concluded a high toxicity to bees even considering the numerous limitations in the studies included into the dossier and a number of sub-lethal effects do not taken into account in the framework of risk assessment (Bee Life 2013 Annex II).

Residues using Good Agricultural Practicies (GAPs)

Residues in fruits and other matrices following an application of sulfoxaflor under GAP are used for the estimation of MRLs. The FAO has summarised the residues in different crops collected from studies carried out worldwide. Highest residue (HR) values in citrus fruits was 0.44 mg kg⁻¹, in pomme fruits 0.26 mg kg⁻¹, in stone fruits 1.2 mg kg⁻¹, in grapes 1.6 mg kg⁻¹, and strawberries 0.21 mg kg⁻¹. On fruiting vegetables and cucurbits the HR value is 0.27 mg kg⁻¹. Leafy vegetables HR value is 2.9 mg kg⁻¹ sulfoxaflor. (FAO 2011)

Analytical sensibility

Currently there is the possibility to detect sulfoxaflor at a level of detection (LOD) of ranging from 0.2 to 0.6 µg kg⁻¹, and LOQs from 0.7 to 2.0 µg kg⁻¹ (Xu et al. 2012). These are more than 100 times lower than the MRLs proposed by the European Commission in (MRL 0.05 mg kg⁻¹).

Annexe 1. Results of acute toxicity tests (laboratory)

Test species	Test substance	Test system / Duration	Endpoint	Reference
Honey bee	Sulfoxaflor (technical)	Lab. acute oral, 48h	LD ₅₀ oral = 0.146 µg a.s./bee	Bergfield, A. 2007a IIA 8.7.1 /01
Honey bee	Sulfoxaflor (technical)	Lab. acute contact, 72h	LD ₅₀ contact = 0.379 µg a.s./bee	Bergfield, A. 2007b IIA 8.7.2/01
Honey bee	GF-2626	Lab. acute oral, 48h	LD ₅₀ oral = 0.539 µg GF-2626/bee LD ₅₀ oral = 0.065 µg Sulfoxaflor/bee	Vinall, S. 2010a IIIA1 10.4.2.1 /01
Honey bee	GF-2626	Lab. acute contact, 48h	LD ₅₀ contact = 2.356 µg GF-2626/bee LD ₅₀ contact = 0.283 µg Sulfoxaflor/bee	Vinall, S. 2010b IIIA1 10.4.2.2 /01
Honey bee	GF-2372	Lab. acute oral, 48h	LD ₅₀ oral = 0.153 µg GF-2372/bee LD ₅₀ oral = 0.075 µg Sulfoxaflor/bee	Vinall, S. 2010 IIIA2 10.4.2.1 /01
Honey bee	GF-2372	Lab. acute contact, 48h	LD ₅₀ contact = 0.224 µg Sulfoxaflor/bee	Vinall, S. 2009 IIIA2 10.4.2.2 /01
Bumble bee	GF-2032	Lab. acute oral and contact, 72h	LD ₅₀ oral = 0.027 µg XDE/208/bee LD ₅₀ contact = 7.554 µg Sulfoxaflor/bee	Vinall, S. 2009 IIIA1 10.4.2.1 /02
Honey bee	X11719474	Lab. acute oral, 48h	LD ₅₀ oral >100 µg./bee*	Vinall, S. 2009 IIA 8.7.1 /02
Honey bee	X11519540	Lab. acute oral, 48h	LD ₅₀ oral >91.2 µg./bee*	Vinall, S. 2010a IIA 8.7.1 /03
Honey bee	X11579457	Lab. acute oral, 48h	LD ₅₀ oral = 45.7 µg./bee	Vinall, S. 2010b IIA 8.7.1 /04
Honey bee	X11721061	Lab. acute oral, 48h	LD ₅₀ oral >103.5 µg./bee*	Vinall, S. 2010c IIA 8.7.1 /05
Honey bee	GF-2032	Laboratory foliar residue toxicity test	No significant adverse effects to bees when exposed to foliar residues of GF-2032 treated 3, 6 or 24 hours previously at 200 g Sulfoxaflor/ha.	Lee, B. 2008 IIIA1 10.4.3/01
Honey bee	GF-2372	Laboratory foliar residue toxicity test	No significant adverse effects to bees when exposed to foliar residues of GF-2372 treated 3, 6 or 24 hours previously at 100 and 200 g Sulfoxaflor/ha.	Bergfield, A. 2009 IIIA2 10.4.3/01

References

- Babcock JM, Gerwick CB, Huang JX, Loso MR, Nakamura G, Nolting SP, et al. Biological characterization of sulfoxaflor, a novel insecticide. Pest management science. 2011;67: 328–334.
- Cutler P, Slater R, Edmunds AJF, Maienfisch P, Hall RG, Earley FGP, Pittnera T, Pal S, Paul VL, Goodchild J, Blacker M, Hagmann L, Crossthwaite AJ (2013) Investigating the mode of action of sulfoxaflor: a fourth-generation neonicotinoid. Pest Manag Sci 69:607–619
- DoW Agro Sciences (2013) DoW AgroSciences receives US EPA Registration for Sulfoxaflor EFSA. Conclusion on the peer review of the pesticide risk assessment of the active substance sulfoxaflor. EFSA Journal. 2014;12: 3692 [170 pp.]. doi:doi:10.2903/j.efsa.2014.3692
- Oliveira EE, Schleicher S, Büschges A, Schmidt J, Kloppenburg P, Salgado VL. Desensitization of nicotinic acetylcholine receptors in central nervous system neurons of the stick insect (*Carausius morosus*) by imidacloprid and sulfoximine insecticides. Insect Biochemistry and Molecular Biology. 2011;41: 872–880. doi:10.1016/j.ibmb.2011.08.001
- Satchivi NM, Schmitzer PR. Synergistic herbicide/insecticide composition containing certain pyridine carboxylic acids and certain insecticides. USA; US20110207606, 2011.
- Shao X, Swenson TL, Casida JE (2013b) Cycloxyprodifopid insecticide: nico- tinic acetylcholine receptor binding site and metabolism. J Agric Food Chem 61:7883–7888. doi:10.1021/jf4030695
- Sparks TC, DeBoer GJ, Wang NX, Hasler JM, Loso MR, Watson GB (2012) Differential metabolism of sulfoximine and neonicotinoid insecticides by *Drosophila melanogaster* monooxygenase CYP6G1. Pest Biochem Physiol 103:159–165
- Sparks TC, Watson GB, Loso MR, Geng C, Babcock JM, Thomas JD (2013) Sulfoxaflor and the sulfoximine insecticides: chemistry, mode of action and basis for efficacy on resistant insects. Pest Biochem Physiol 107:1–7
- US Environmental Protection Agency (USEPA) (2013) Registration of the new active ingredient sulfoxaflor for use on multiple commodities, turgrass and ornamentals. United States Environmental Protection Agency <http://www.regulations.gov/#!documentDetail;D=EPA-HQ-OPP-2010-0889-0396>. Accessed 7 Mar 2014
- Watson GB, Loso MR, Babcock JM, Hasler JM, Letherer TJ, Young CD, et al. Novel nicotinic action of the sulfoximine insecticide sulfoxaflor. Insect biochemistry and molecular biology. 2011;41: 432–439.
- Xu J, Dong F, Liu X, Li J, Li Y, Shan W, et al. Determination of sulfoxaflor residues in vegetables, fruits and soil using ultra-performance liquid chromatography/tandem mass spectrometry. Anal Methods. 2012;4: 4019–4024. doi:10.1039/C2AY25782C