

**WARNING KEEP OUT OF REACH OF CHILDREN**



**For the control or suppression of the major seed borne and soil borne diseases in cereals, maize and sweetcorn.**

**GROUP 3 & 4 FUNGICIDE**

ACTIVE INGREDIENT:

25g/litre ipconazole and 20g/litre metalaxyl in the form of a microemulsion.



**AGRECOVERY**  
*Rural Recycling Programme*

Approved pursuant to HSNO Act 1996, EPA Approval Code HSR101127

See <http://www.epa.govt.nz> for approval controls.

Registered pursuant to the ACVM Act 1997, No. P009285

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**NET  
CONTENTS:**

**20**

**LITRES**

**Arysta LifeScience NZ**

**PO Box 51584, Pakuranga**

**Auckland, New Zealand**

**0800 100 325**

**WWW.ARYSTALIFESCIENCE.NZ**



**Arysta**  
LifeScience

**PLEASE READ THE LABEL COMPLETELY BEFORE USE**

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and conditions of registration.

This product **MUST** be under the control of an Approved Handler during use.

**HSNO CLASSIFICATION: 6.9B, 9.1C****WARNING**

May cause damage to organs through prolonged or repeated exposure.

Harmful to aquatic life with long lasting effects.

**PRECAUTIONS**

Store in original container tightly closed, away from foodstuffs.

When handling product and when cleaning seed treatment equipment wear cotton overalls, chemical resistant apron, elbow length chemical resistant gloves, waterproof boots and a face shield or goggles.

When handling treated seed wear cotton overalls and elbow length chemical resistant gloves.

Avoid contact with skin and eyes, and inhalation of spray mist.

DO NOT eat, drink or smoke while using.

Wash hands and face before eating, drinking or using the toilet and after work.

Apply using properly maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

DO NOT contaminate any aquatic environment with chemical mixture or empty container.

**CONTAINER DISPOSAL:** Ensure container is completely empty. Triple rinse the empty container with soap and water. Recycle through the Agrecovery programme. If this is not possible, bury in an approved landfill or other approved facility.

Do not dispose of this product down drains or sewers.

**PRODUCT DISPOSAL:** Ensure unwanted product is used by another operator strictly in accordance with label directions or at an approved landfill or other approved facility.

**FIRST AID:** If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

**SPILLAGE:** Wear appropriate protective clothing and prevent material from entering waterways. Contain liquid spills and absorb with inert material. Dispose of waste safely in a suitable landfill.

**RESISTANCE MANAGEMENT:** RANCONA® DIMENSION fungicide contains ipconazole, a fungicide from the triazole group of fungicides [GROUP3 FUNGICIDE] with the DMI (Demethylation Inhibitors) mode of action, and metalaxyl, a fungicide from the phenylamide group of fungicides [GROUP4 FUNGICIDE]. Some naturally occurring fungi resistant to these groups of fungicides may exist through normal genetic variability in any fungal population. Resistance to this product could develop from excessive use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies that exist for any fungus listed on the label. Refer to [www.resistance.nzpps.org/](http://www.resistance.nzpps.org/) for further information.

**DIRECTIONS FOR USE:**

**SEED QUALITY:** Seed treatment should not be used on seed that is more than 12% moisture content or on sprouted, damaged or severely pinched seed, or seed of poor viability. If in doubt have a germination test carried out on seed before treatment to ensure that it is of acceptable standard.

**DISEASES CONTROLLED AND USE RATE:**

Crop	Disease	Rate	Comments
Cereals (Wheat, Barley, Oats, Triticale, Ryecorn)	Loose Smut, Covered Smut, Bunt, Seed borne seed and seedling rot diseases (including Rhizoctonia, Pythium, Fusarium spp., M. nivale)	Apply 3.2L/tonne seed	RANCONA DIMENSION can be used neat or diluted with water to achieve optimum coverage. If diluting with water, as a guide add 2-5L water/tonne seed.
Maize and Sweetcorn	Head smut, Seed borne seed and seedling rot diseases (including Rhizoctonia, Pythium and Fusarium spp.)	Apply 3.2L/tonne seed	Ensure thorough and complete coverage of the seed. If using water, the mixture should be gently stirred regularly.

**METHOD OF TREATMENT AND DILUTION RATE:** RANCONA DIMENSION may be applied to seed undiluted or may be mixed with water if required to achieve optimum coverage of the grain. This requirement will depend on the equipment being used. Use only through equipment designed to apply suspension concentrate or slurry seed treatments to grain. Uniform seed treatment is important to ensure the best disease protection and optimum performance.

Diluted product should be agitated before and during application. Do not allow diluted product to stand for long periods e.g. overnight, without agitating thoroughly before continuing application.

RANCONA DIMENSION may be applied to seed at the same time as other seed treatments (such as insecticides). Dilute Rancoana Dimension and any other seed treatment in an appropriate amount of water to achieve thorough coverage. Do not pre mix individual product concentrates

RANCONA DIMENSION must not be applied at rates exceeding 320ml/100kg seed.

Treated seeds must not be sowed at rates exceeding 200kg/ha.

RANCONA DIMENSION must be applied to seeds in a purpose-built seed coating machine.

**STORAGE OF TREATED SEED:** If seed is not used immediately it should be stored in a dry, well ventilated place. Although RANCONA DIMENSION has no effect on the viability of treated seed, subsequent germination may be affected by seed quality or by poor storage conditions such as high moisture and/or high temperatures. No liability can be accepted for the performance of stored treated seed.

**WITHHOLDING PERIOD:**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

DO NOT feed treated crops to animals for 10 weeks after sowing.

**Containers of treated seed must be labelled with the following:**

- Treated seed must not be fed to humans, animals or birds.
- Stock must not graze maize, sweetcorn or cereals for 10 weeks after planting.
- Treated seeds must be completely covered by soil when sown and excess treated seed must not be left in areas accessible to birds.

RANCONA is a registered trademark of MacDermid Agricultural Solutions Inc.

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

Registered to ETEC Crop Solutions Limited, an Arysta LifeScience Group Company. 0800 100 325

**EXCLUSION OF LIABILITY:**

**Please read these conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened.**

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, ETEC Crop Solutions Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Proper Shipping Name	Hazard Class	UN No.	Hazchem	Packing Group
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (ipconazole 2.5%)	9	3082	2Z	III

**In a transport emergency dial 111,  
Police or Fire Brigade  
For specialist advice  
in an emergency only,  
call 0800 CHEMCALL (0800 243 622)**

<b>BATCH NO</b>
<b>DOM</b>



AR185/15L

9 421022 425984