

Table 5. 2019 American Geriatrics Society Beers Criteria® for Potentially Clinically Important Drug-Drug Interactions That Should Be Avoided in Older Adults

Object Drug and Class	Interacting Drug and Class	Risk Rationale	Recommendation	Quality of Evidence	Strength of Recommendation
RAS inhibitor (ACEIs, ARBs, aliskiren) or potassium-sparing diuretics (amiloride, triamterene)	Another RAS inhibitor (ACEIs, ARBs, aliskiren)	Increased risk of hyperkalemia	Avoid routine use in those with chronic kidney disease stage 3a or higher	Moderate	Strong
Opioids	Benzodiazepines	Increased risk of overdose	Avoid	Moderate	Strong
Opioids	Gabapentin, pregabalin	Increased risk of severe sedation-related adverse events, including respiratory depression and death	Avoid; exceptions are when transitioning from opioid therapy to gabapentin or pregabalin, or when using gabapentinoids to reduce opioid dose, although caution should be used in all circumstances.	Moderate	Strong
Anticholinergic	Anticholinergic	Increased risk of cognitive decline	Avoid; minimize number of anticholinergic drugs (Table 7)	Moderate	Strong
Antidepressants (TCAs, SSRIs, and SNRIs) Antipsychotics Antiepileptics Benzodiazepines and nonbenzodiazepine, benzodiazepine receptor agonist hypnotics (ie, "Z-drugs") Opioids	Any combination of three or more of these CNS-active drugs ^a	Increased risk of falls (all) and of fracture (benzodiazepines and nonbenzodiazepine, benzodiazepine receptor agonist hypnotics)	Avoid total of three or more CNS-active drugs ^a ; minimize number of CNS-active drugs	Combinations including benzodiazepines and nonbenzodiazepine, benzodiazepine receptor agonist hypnotics or opioids: high All other combinations: moderate	Strong
Corticosteroids, oral or parenteral	NSAIDs	Increased risk of peptic ulcer disease or gastrointestinal bleeding	Avoid; if not possible, provide gastrointestinal protection	Moderate	Strong
Lithium	ACEIs	Increased risk of lithium toxicity	Avoid; monitor lithium concentrations	Moderate	Strong
Lithium	Loop diuretics	Increased risk of lithium toxicity	Avoid; monitor lithium concentrations	Moderate	Strong
Peripheral α-1 blockers	Loop diuretics	Increased risk of urinary incontinence in older women	Avoid in older women, unless conditions warrant both drugs	Moderate	Strong
Phenytoin	Trimethoprim-sulfamethoxazole	Increased risk of phenytoin toxicity	Avoid	Moderate	Strong
Theophylline	Cimetidine	Increased risk of theophylline toxicity	Avoid	Moderate	Strong
Theophylline	Ciprofloxacin	Increased risk of theophylline toxicity	Avoid	Moderate	Strong
Warfarin	Amiodarone	Increased risk of bleeding	Avoid when possible; if used together, monitor INR closely	Moderate	Strong
Warfarin	Ciprofloxacin	Increased risk of bleeding	Avoid when possible; if used together, monitor INR closely	Moderate	Strong
Warfarin	Macrolides (excluding azithromycin)	Increased risk of bleeding	Avoid when possible; if used together, monitor INR closely	Moderate	Strong
Warfarin	Trimethoprim-sulfamethoxazole	Increased risk of bleeding	Avoid when possible; if used together, monitor INR closely	Moderate	Strong
Warfarin	NSAIDs	Increased risk of bleeding	Avoid when possible; if used together, monitor closely for bleeding	High	Strong

Abbreviations: ACEI, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; CNS, central nervous system; INR, international normalized ratio; NSAID, nonsteroidal anti-inflammatory drug; RAS, renin-angiotensin system; SNRI, serotonin-norepinephrine reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor; TCA, tricyclic antidepressant.

^aCNS-active drugs: antiepileptics; antipsychotics; benzodiazepines; nonbenzodiazepine, benzodiazepine receptor agonist hypnotics; TCAs; SSRIs; SNRIs; and opioids.