

CURFEW[®]

FUNGICIDE

LIGHTS OUT FOR
ONION DOWNY
MILDEW AND
POTATO LATE
BLIGHT

NEWS

- ✓ POWERFUL 24-48 HOUR CURATIVE ACTIVITY POST INFECTION
- ✓ FORWARD PROTECTION UP TO THREE DAYS
- ✓ OUTSTANDING TANK MIX PARTNER FOR LATE BLIGHT & DOWNY MILDEW
- ✓ NO KNOWN RESISTANCE AFTER DECADES OF USE
- ✓ PROVEN UNDER HIGH PRESSURE EUROPEAN CONDITIONS
- ✓ VERY COST EFFECTIVE
- ✓ CLEAN UP DISEASE BEFORE IT GETS UNDERWAY

CURFEW[®]

FUNGICIDE

TECHNOTE



USE CURFEW AS A TANK MIX PARTNER AND CLAMP DOWN HARD ON DISEASE

Curfew contains cymoxanil (450g/kg) in a water dispersible granule. With both translaminar and penetrant qualities **Curfew** provides superior curative activity against late blight and downy mildew. Trials confirm cymoxanil in **Curfew** has strong curative action for 24 – 48 hours after an infection has taken place plus up to three days forward cover.



CURFEW REINFORCES DISEASE CONTROL

Curfew is the perfect attacking ally to complement a defensive disease program. With proven kick back curative activity **Curfew** goes on the offensive with punch and flexibility as a tank mix partner with contact protectant fungicides such as mancozeb.



CURFEW A TRUSTED RESISTANCE FIGHTER

Resistance to phenylamide fungicides is well documented and many new single site fungicides are also susceptible. Cymoxanil is a member of the acetamid group, uniquely different to any currently available fungicides in New Zealand. With multi-site activity, resistance to **Curfew** is unlikely and yet to be documented.



KNOWING WHEN TO IMPOSE CURFEW

Don't let things get out of control, enforce a **Curfew** as soon as a disease event occurs and before the first symptoms of unrest (disease) become visible. The optimum approach is to use a disease monitoring service and apply within 24 – 48 hours of an infection taking place but before the disease is apparent in the crop. If disease pressure is high don't back down, re-impose a **Curfew**, look at two applications of **Curfew** back to back at close intervals (5 days apart) and curb unwanted behavior (disease spread).

Crop	Disease	Critical Comments
Onions and Garlic	Downy Mildew	Apply 250 – 350g / ha. Use as a tank mix partner with a protectant fungicide. Begin spraying at the 2 true leaf stage and continue on a 7 to 10 day schedule, using Curfew within 48 hours of an infection period. Do not use more than 3 times in succession and do not apply more than 5 times per season.
Potatoes	Late Blight	Apply 220 - 250g / ha as a tank mix partner with a protectant fungicide at 7 to 10 day intervals. Use Curfew within 48 hours of an infection period and use alternative fungicides at other times. Do not use more than 3 times in succession and do not apply more than 5 times per season. Use the higher rate, and two applications 5 to 7 days apart under extreme infection conditions, i.e. warm with high humidity.



ADJUVANTS

Good coverage is essential for optimum disease control. When applying **Curfew** in 300 L of water per hectare or less always use Du-Wett[®] or Du-Wett[®] Rainmaster low water volume adjuvant systems to achieve superior levels of coverage. In water volumes above 300 L Bond[®] Xtra or Actiwett[®] are recommended. If irrigation is scheduled or rainfall expected soon after application use Du-Wett Rainmaster (< 300 L / Ha) or Bond Xtra (> 300 L / Ha) for wash off protection.

Pack size 2.5kg.

WITHHOLDING PERIOD 14 DAYS

Etec Crop Solutions Ltd.

PO Box 51-584, Pakuranga, Auckland, New Zealand

NZ Freephone: 0800 100 325. Telephone: +64 9 574 5401

For more information on Etec Crop Solution products visit our website at

www.etec.co.nz



Registered pursuant to the ACVM Act 1997, No. P8010