

ManKocideTM

Fungicide

Bactericide

Dual Action for Double Protection

- ✓ Copper + Mancozeb in a superior DF formulation
- ✓ Broad spectrum bactericide/fungicide
- ✓ Excellent mixing/spraying and compatibility

Benefits

- Synergies from the combination
- Broader disease spectrum
- Control of Copper tolerant bacteria
- Reduced phytotoxicity in grapes
- Convenience to the grower
- Less packaging



www.etec.co.nz

Etec
CROP SOLUTIONS


16000916



30% metallic Copper + 15% mancozeb

ManKocide® is a convenient broad spectrum fungicide/bactericide, formulated to provide maximum protection against fungal and bacterial diseases.

- **ManKocide measures easily**
- **Pours with minimal dust**
- **Mixes readily in the spray tank**



ManKocide protects potatoes from late blight, using less Mancozeb (23%) and less Copper (63%) than the standards.

Dry flowable formulation

Broad spectrum disease protection

Resistance management tool

Product wets instantly

Tank-mix compatibility

ManKocide can be mixed with foliar applied fungicides, insecticides and most foliar nutrients. The formulation has two active ingredients eliminating the need to apply two separate formulations in the spray tank helping to reduce potential mixing problems.

Do not mix with strongly acidic or alkaline material.

For further information contact: Etec Crop Solutions Ltd.
PO Box 51584, Pakuranga 2140, Auckland, New Zealand.

Phone: 0800 100 325 www.etec.co.nz

© Copyright Etec Crop Solutions Limited 2016, all rights reserved. All directions or use, recommendations, use rates, compatibilities and other technical information is proprietary to Etec Crop Solutions. ® Registered Trademarks ®MANKOCIDE is a registered trademark of Kocide LLC. Registered pursuant to the ACVM Act 1997, No.P005674 MANK1016

WHY **ManKocide**?

A quality formulated pre mix of mancozeb and copper (Kocide®) that has proven to be more effective than either product alone or tank mixed, and equal to over night tank mixes.

GETTING THE BEST FROM **ManKocide**

Spray coverage is essential for consistent disease control. Apply ManKocide with Actiwett® in dilute spray volumes or with Du-Wett® in concentrate spray volumes to achieve complete coverage without undue run-off.

If a rainfast adjuvant is required, use Bond Xtra® or Du-Wett WeatherMax® instead.

DO NOT apply less than the recommended minimum rate of ManKocide.

RECOMMENDATIONS

Grapes – Black spot, downy mildew

Apply 200 - 250 g/100 litres water (minimum 2.5 kg/ha) at 10 – 14 day spray intervals during the pre-bloom and post-bloom periods.

Potatoes – Late blight, early blight

Apply 2.5 - 4 kg per ha at 7 - 10 day intervals. Use the higher rate when conditions favour disease development.

Pumpkins, Squash – Angular leaf spot, downy mildew

Apply 2 - 2.5 kg ha at 7 - 10 day intervals commencing when plants begin to run.

Tomatoes - Late blight, early blight, bacterial speck, bacterial spot

Apply 250 - 400 g/100 litres of water/ha (minimum 2.5 kg/ha) – at 7 - 10 day intervals. Repeat applications after 13 mm rain or irrigation.

Withholding periods

Cucurbits and tomatoes – 1 day, all Other crops – 14 days

ManKocide® DF may be measured volumetrically.

Pour ManKocide® DF slowly into a dry measuring cylinder or jug.

Approx volumetric measure 1kg = 950mL

NOTE: Measurement by weight is the most accurate.

15kg Pack