

**KEEP OUT OF REACH OF CHILDREN
DANGER**

Capetec

Fungicide for the control of diseases in fruit, vegetable, ornamental and green house crops and for use on turf, ornamentals and certain soil borne diseases.

Read the Label Carefully before use

Active Ingredient:
800 g/kg captan in the form of a water dispersible granule.

GROUP M4 FUNGICIDE

Registered pursuant to the ACVM Act 1997, P8921.
See: www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996. Approval Code HSR000503.

Distributed By:

Etec Crop Solutions Limited
PO Box 51584, Pakuranga
Auckland, New Zealand
Phone: 0800 100 325
www.etec.co.nz

NET
CONTENTS:

10
Kg

Etec
CROP SOLUTIONS

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and conditions of registration.

This product **MUST** be under the control of an Approved Handler during use.



HSNO Classifications: 6.3B, 6.5B, 6.7B, 8.3A, 9.1A

DANGER

May cause skin irritation.
May cause an allergic skin reaction.
Suspected of causing cancer.
Causes serious eye damage.
Very toxic to aquatic life.

PRECAUTIONS

Store in original container tightly closed, away from foodstuffs.
When mixing or applying, wear protective clothing, gloves and goggles.
Avoid contact with skin and eyes, and inhalation of spray mist.
Wash splashes of concentrate from skin or eyes immediately.
If clothing becomes contaminated with product, remove clothing immediately.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
Wash protective clothing after use.
Apply with well maintained and calibrated spray equipment.
Records of use **MUST** be kept.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container.

CONTAINER DISPOSAL: Ensure container is completely empty. Crush and bury in suitable landfill.

PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

SPILLAGE

Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. This product is classified under HSNO as 9.1A, and storage of certain quantities may require bunding and signage. See the HAZNOTE for further information.

SAFETY DATA SHEET

If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.etec.co.nz

24 HOUR EMERGENCY RESPONSE
0800 CHEM CALL 0800 243 622

BATCH NO

DOM



EL105/10KG

9 421022 424987

GENERAL INFORMATION:

Capetec is a broad spectrum contact, protectant fungicide from the cyclic imide chemical group. Capetec has no systemic activity so complete coverage of the target is essential for best results.

COMPATIBILITY:

Capetec is compatible with most commonly used pesticides. Do not apply with highly alkaline products such as Bordeaux mixture or lime Sulphur. Do not mix with oils or emulsifiable concentrates. Capetec mixed with Sulphur and used on the apple varieties Kidds Orange, Delicious and other 'Sulphur shy' varieties, may cause leaf spotting and leaf drop may occur.

MIXING:

Partly fill the spray tank. Turn on agitation and while continuing to fill the tank, add the required amount of Capetec directly into the tank. Maintain agitation during spraying and agitate thoroughly after a stoppage. Add any tank mixing partners just before tank filling is complete.

Knapsack Sprayer Rates:

Hydraulic type knapsack: 13 grams per 10 litres of water.
 Motorised airblast knapsack: 40 grams per 10 litres of water. Use the middle settings on the restrictor and operate at sufficient revs to produce small droplets.

DIRECTIONS FOR USE:

It is an offence to use this product on animals

The rates suggested are for high volume spraying to the point of run-off. For concentrate spraying, adjust dilution rate accordingly. Use the highest recommended rate and/or the shortest spray interval when disease pressure is high. Ensure thorough coverage of all parts of treated plants.

RECOMMENDATIONS

FRUIT CROPS			
CROP	DISEASE	RATE	REMARKS
Apples & Pears	Black spot Glomerella Ripe spot Sooty blotch	75-125 g / 100 litres of water	For blackspot control apply the high rate at 7 day intervals from green tip to petal fall. The rate may be reduced and the interval may be increased to 10-14 days, from petal fall to harvest. To control glomerella, ripe spot and sooty blotch, use the high rate from January through to harvest. CAPETEC may increase the risk of russet on russet sensitive varieties. Use alternative fungicides on these varieties from full bloom to second cover. DO NOT APPLY TO ANJOU PEARS.
Blueberries	Stem blight Leaf spot Botrytis	125 g / 100 litres of water	Apply just prior to flowering, at 10% flowering and at petal fall. Continue to spray at 10-14 day intervals until harvest.
Boysenberries Loganberries Other Brambles	Cane spot	125 g / 100 litres of water	Apply at petal fall and repeat at 10-14 day intervals until harvest.
Citrus	Brown rot	125-240 g / 100 litres of water	Apply as a skirt spray before winter. Wet trees thoroughly to a height of 1 metre. Repeat 10 weeks later. In Autumn, treat whole tree and repeat 4-6 weeks later.
Black & Red Currants	Leaf spot Botrytis	125 g / 100 litres of water	Apply at 10-14 day intervals from green tip until harvest.
Gooseberries	Leaf spot Botrytis	125 g / 100 litres of water	Apply just prior to flowering, and repeat at petal fall, preharvest and postharvest. To control Botrytis flower rot, apply at 5-7 day intervals over flowering when infection conditions are favourable for disease.
Grapes	Botrytis Dead arm Anthracnose	125 g / 100 litres of water	Apply at 14 day intervals from the time when shoots are 50-80 mm long.
	Downy mildew	90 g / 100 litres of water	Apply at the start of flowering. Use an alternative botryticide over the remainder of the flowering period or use CAPETEC at 125 g / 100L for Botrytis control. Continue applications at the 90 g rate at 14 day intervals after flowering.
Raspberries	Cane spot Spur blight Botrytis	125 g / 100 litres of water	Main crop: apply at 10-14 day intervals from green tip until harvest. Autumn crop: Apply at 10-14 day intervals from November until harvest. Note: Thorough coverage of plants is essential.
Stone fruit	Brown rot Scab Shothole	125 g / 100 litres of water	Apply from bud movement until bloom at 7 day intervals, then at 3-4 day intervals until petal fall. Continue applications at 14 day intervals until 3 weeks before harvest and then apply at 7 day intervals until harvest particularly in wet conditions.
Strawberries	Leaf spot Botrytis	3 kg/ha or 125g / 100 litres of water	Apply as required to control leaf spot. For botrytis control, apply 3 kg/ha at 7 day intervals from the start of flowering and between pickings if warm wet conditions prevail.

VEGETABLE CROPS			
CROP	DISEASE	RATE	REMARKS
Peas	Root rot	7-9 kg / ha	Apply as a broadcast spray before planting and incorporate into the top 100mm of soil.
Beans	Damping-off		
Carrots	Ring spot Cercospora blight	3 kg / ha	Apply as required
Cucurbits	Anthracnose Downy mildew	1.5 – 3 kg / ha	Apply as required in sufficient water to obtain complete coverage
Lettuce	Leaf spot	4.5 kg / ha	Apply at 10-14 day intervals from the rosette stage.
Potato Seed	Seed rot Damping off	125 g / 100 litres of water	Dip seed potatoes within 6 hours of cutting.

AMENITY TURF AND ORNAMENTALS			
CROP	DISEASE	RATE	REMARKS
Lawns	Brown patch Leaf spot Damping off	6 kg / ha (6 grams / 10 square metres)	Apply at 7-10 day intervals when diseases are prevalent in summer and autumn. Apply as a soil drench in sufficient water to penetrate into the soil.
Ornamentals	Rose black spot Carnation leaf spot Chrysanthemum spot	125 g / 100 litres of water	Apply as required.
Soil and glasshouse soil drench	Damping-off	125 g/100 litres Apply to 100 square metres	Apply and work into the top 100mm of soil before planting

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS:

Boysenberries	14 days
Blueberries	1 day
Carrots	14 days
Citrus	28 days
Cucurbits	3 days
Currants	1 day
Gooseberries	1 day
Grapes	21 days
Lettuce	7 days
Loganberries and other Brambles	14 days
Strawberries	1 day
Pip fruit	14 days
Raspberries	1 day
Stonefruit	7 days

For export crops – consult with your exporter for advice about any residue restrictions that may apply to the intended importing country

Exclusion of Liability: Please Read

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Etec Crop Solutions Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered pursuant to ACVM Act 1997, No. P8258
 HSNO Approval Code HSR000503
 See www.foodsafety.govt.nz for registration conditions.

**UN No: 3077,
 ENVIRONMENTALLY
 HAZARDOUS SUBSTANCE
 N.O.S. (Contains 80% CAPTAN)
 MARINE POLLUTANT
 Packing Group III, DG 9**

**In a transport emergency
 dial 111, Police or Fire Brigade
 For specialist advice in an emergency only,
 call 0800 CHEMCALL (0800 243 622)**

