

**HARMFUL. ECOTOXIC
KEEP OUT OF REACH OF CHILDREN**



Potato Dormancy Enhancer and Sprout Inhibitor

READ THE LABEL CAREFULLY BEFORE USE



AGRECOVERY

Active Ingredient:
960 g/ Litre 1,4-dimethylnaphthalene
in the form of a technical concentrate (DMN).

Registrant:
Etec Crop Solutions Limited
PO Box 51584, Pakuranga 2140
Auckland, New Zealand
Phone: 0800 100 325
www.etec.co.nz



NET
CONTENTS:

10
Litres

Etec
CROP SOLUTIONS

180mm High x 210mm Wide



CMYK

HSNO Class: 6.1E, 6.3A, 6.4A, 9.1A

Approved Handler Applies:

This product must be under the personal control of an approved handler or secured

ENVIRONMENTAL WARNING: ECOTOXIC - Very toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. HARMFUL to the soil environment and to terrestrial vertebrates.

HEALTH WARNING: HARMFUL - may be harmful if swallowed, inhaled or absorbed through the skin. May cause skin and eye irritation. Avoid contact with the skin and eyes, avoid inhalation.

PRODUCT DISPOSAL: Ensure unwanted product is used by another operator strictly in accordance with label directions.

CONTAINER DISPOSAL: Ensure container is completely empty. Triple rinse and submit clean empty container for recycling through Agrecovery. If this is not possible, burn if permitted and circumstances, especially wind direction, allow; otherwise bury in landfill.

HANDLING PRECAUTIONS: When mixing or applying, wear protective clothing, gloves and goggles. Do not eat, drink or smoke while using. Wash hands and face before meals and after work. Wash protective clothing after work. Store in original container away from feed, seeds and foodstuffs. Flush spray equipment thoroughly with clean water after use.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. This product is classified under HSNO as 9.1A, and storage of certain quantities may require bunding and signage. See the HAZNOTE for further information. When stored appropriately this product should show no significant degradation for a minimum of 2 years.

FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the NATIONAL POISONS CENTRE 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT SCRUB THE SKIN.

PRODUCT STORAGE CONDITIONS: 1,4SIGHT™ will freeze at temperatures below 5°C. Store in a warm (15-20°C) store.

GENERAL INFORMATION:

The active ingredient (1,4-dimethylnaphthalene – (1,4-DMN)) occurs naturally in potatoes. While it remains present at sufficient levels, sprouting is prevented. This dormancy enhancement mode of action is different to treatment with CIPC which burns the sprouts after dormancy has broken. Over time, potatoes that have been harvested or ground stored, lose 1,4-DMN and sprout production starts. Potatoes treated with 1,4SIGHT have less weight loss and remain more turgid than untreated. If sprouting has already started at the time of treatment, the highest rate recommended for 1,4SIGHT may only slow the rate of sprouting but may not stop it, depending on variety. By applying 1,4SIGHT to stored potatoes and maintaining the levels of 1,4-DMN, the natural process for controlling sprouting is maintained.

USE CONDITIONS:

When applied to storage areas 1,4SIGHT is absorbed into the potatoes. The level of 1,4-DMN in the potatoes should not be allowed to drop below 1 ppm. This can be checked by periodic (2-4 weekly) residue analysis of the tubers. In the absence of residue analysis regular inspections should be made and retreatment made at the first signs of sprouting ('peeps'). Target inspections to the areas of the store where it is most difficult to get good vapour treatment penetration. Regular inspections should also be made for rotting potatoes as 1,4SIGHT can mask the odour of rotting potatoes. Open only the minimum space required for ingress when making inspections to minimize loss of vapours from the store.

For optimum sprout control, potatoes should be dug immediately on maturity, moved into storage and treated. Suberization will not be impaired.

Potatoes that have been ground stored before harvest will contain low natural levels of 1,4-DMN and may require more frequent treatment of 1,4SIGHT to establish and maintain base levels.

Best results are in Potato stores where there is minimal air exchange and no fresh air into the store for 24-48 hours after treatment. 1,4SIGHT may leak from storage areas that are not well sealed. Such stores will require more frequent re-treatment to maintain levels of 1,4SIGHT. Areas of potential leakage should be minimised for economic control of sprouting. 1,4SIGHT vapours can pass through wood and most plastics. Do not use these materials to repair leaks.

1,4SIGHT should be applied as an vapour treatment in a cold or hot fog to the air space within the store. Use thermal fogging under cooler temperatures. Application as a spray 'mist' can result in 'spotting' of the potatoes. To assist with penetration into the stack, air vents under the stack are recommended. Where none exist, effective air ducts can be made, by using normal pallets stacked two high and butted open end, to open end. Under stack ducting should be not more than 3 metres apart under the stack.

Best results will be achieved if the storage temperature of the potatoes is maintained at or below 10°C and all other conditions for good long term storage of potatoes are met. There are major differences between variety dormancy properties and sprouting potential. It is important to inspect for the first sign of sprouting at regular intervals. Re-treat if any indications that dormancy has broken.

Do not store Seed Potatoes for immediate planting (5-6 weeks) in any area that shares air space with 1,4SIGHT treated potatoes. For treatment of 'Shipping containers' refer to the product 1,4SHIP®.

RATE OF APPLICATION:

Apply 10 -20 ml / Tonne of Potatoes.

Use the high rate for the initial treatment. The low rate should be used for subsequent re-treatments as required to maintain dormancy unless the store is only partially filled, in which case the high rate should be continued. Up to 30 days Sprout Suppression can be expected from each treatment if storage conditions are favourable. 1,4SIGHT can be used repeatedly until potatoes are sold

METHODS OF APPLICATION:

The following application equipment types are suitable for application:

- Wet fog equipment and Misting equipment feeding into forced aeration systems.
 - These should only be used when in-store temperatures are above 10°C. In colder conditions droplets rather than a vapour are formed. Droplets settle on the potato skins and can mark the tubers. Do not allow any liquid 1,4SIGHT to contact the surface of the potato. Do not allow any liquid 1,4SIGHT to contact plastic in the store. Operate the ventilation at as low a volume as reasonable.
 - Thermal Fogging equipment capable of generating a dry fog, is the preferred method of application. Do not allow any liquid to contact the surface of the potatoes. Do not allow any liquid to contact plastic in the store. Operate the ventilation at as low a volume as reasonable.
- After application the normal store ventilation system should be set to recirculate only and should remain on this setting for 24 - 48 hours to allow the vapours to fill the air space and be absorbed by the tubers. Maintain a gentle air circulation within the store during this 24 - 48 hour period to aid movement of the vapours through-out the whole stack.

After 48 hours resume normal ventilation procedures

- Monitor the potatoes regularly at least once a week. If not using analysis to determine periodic DMN levels, monitor potatoes more frequently, particularly close to doors etc where leakage could be expected, and, at the first sign of any eye movement, re-treat the whole store with 10 ml / tonne 1,4SIGHT in the same manner as described above with reference to air movement and ventilation. Resume monitoring after re-treatment.

RECOMMENDATIONS

GROUND STORED POTATOES:

Where potatoes have been stored in the ground for 4 or more weeks after maturity (or top desiccation) treatment should be made as soon as is practical after placement in the store. Suberization will not be impaired.

FRESHLY HARVESTED POTATOES:

For potatoes that have been harvested or desiccated and then harvested at maturity, treatment may be made as soon as the store is filled or harvest is completed within 3 or 4 weeks. Suberization will not be impaired so double handling and grading is not required before treatment. Treatment should be carried out as soon as practical after placement in long term storage.

TREATMENT OF PACKAGING:

It is possible to treat cardboard and multiwall paper packaging with 1,4SIGHT just before packing to maintain dormancy and reduce sprouting of the potatoes within the packaging.

Apply a measured dose that corresponds to 0.02 ml / kg of packed Potatoes to an area that is not in direct contact with Potatoes to avoid skin burning.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Etec Crop Solutions Ltd Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

© Copyright Etec Crop Solutions Limited 2013 all rights reserved. All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to Etec Crop Solutions.

Batch No.

DOM

EL710

24 Hour Emergency Response
0800 CHEM CALL
0800 243 622



TM is a trademark of Etec Crop Solutions
See www.foodsafety.govt.nz for registration conditions
Registered pursuant to the ACVM Act 1997, P5791

180mm High x 210mm Wide



CMYK