

KEEP OUT OF REACH OF CHILDREN
ECOTOXIC

SupermanZ™

**Fungicide for the control of a range of diseases
in a range of crops.**

Registered pursuant to the ACVM Act 1997 No.P8220
HSNO Approval Code HSR000462
See www.foodsafety.govt.nz for registration conditions

Active Ingredient: 455g/litre mancozeb in
the form of a suspension concentrate.

Group M3 Herbicide

Read the Label Carefully Before Use.



Distributed by:
Etec Crop Solutions Limited
PO Box 51584, Pakuranga
Auckland, New Zealand
Phone: 0800 100 325
www.etec.co.nz



NET
CONTENTS:
10
Litres

FRONT Label 150x150

HSNO CLASSIFICATIONS: 6.4A, 6.5A, 6.5B, 6.9B, 9.1A.

APPROVED HANDLER: This product must be under the control of an approved handler during use. Use of this product in any manner contrary to this label may be an offence under the HSNO Act
HARMFUL: May cause eye irritation. May cause sensitisation from prolonged skin contact. May cause liver damage from repeated oral exposure at high doses.

ECOTOXIC: Very toxic to aquatic organisms. Avoid contamination of water supply with product or empty container.

PRECAUTIONS: Avoid contact with skin and eyes. Avoid breathing dust or spray mist. Do not eat, drink or smoke while using. When mixing or applying wear appropriate protective clothing such as waterproof gloves, hat, overalls, safety goggles and footwear. Wash hands and face before meals and after work. Wash splashes of concentrate from skin and eyes immediately.

Store in original container, tightly closed and under lock and key. Do not store with Class 1,2,3,2,4 or 5 substances or feed, seeds or foodstuffs. Stores containing more than 100 litres of this product require secondary containment and are subject to signage. When stored appropriately, this product should show no significant degradation for 6 months from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

PRODUCT DISPOSAL: Ensure unwanted product is used by another operator strictly in accordance with label directions.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Recycle through the Agrecovery Programme. If this is not possible, burn if permitted and circumstances, especially wind direction permit, otherwise bury in landfill.

CROP	DISEASE	RATE	REMARKS
FRUIT Apples and Pears	Black spot, ripe spot, glomerella, sooty blotch	210-265ml per 100litres	Apply at 10-14day intervals from 3rd cover. Early season use during the main russet sensitive period is not recommended. Use the higher rates where glomerella is a problem.
	Anthracoese (black spot), downy mildew, dead arm disease	350ml per 100litres	
Passionfruit	Brown spot, septoria spot	350ml per 100litres	Apply in an alternating programme with copper sprays on a 14-21 day schedule.
Peaches, Nectarines	Rust	265ml per 100litres	Apply in early November and repeat 2 or 3 times at 14-21 day intervals. To avoid spray deposits on fruit, do not apply within 6 weeks of harvest.
Berry fruit: Blackberries	Rust, cane spot, dryberry	265 -350ml per 100litres	Apply at 10-14 day intervals from shoot development up to petal fall and again after harvest.
Strawberries	Mycosphaerella leaf spot	350ml per 100litres	Apply in mid-June and repeat at 10-14 day intervals or closer in wet conditions.
Other berryfruit	Leaf spot, rust, downy mildew (dry berry)	350ml per 100litres	Apply at greentip and repeat every 10-14 days.
VEGETABLES Cabbage, Celery	Downy mildew, Septoria leaf spot	265-350ml per 100litres 3.2 - 4.4 L/ha	Apply at 3-7day intervals in the seed bed and 7-10 day intervals in the field.
Carrots	Cercospora leaf spot, downy mildew, rust, pleospora spot, anthracnose	265 - 350 ml/100 litres 3.2 - 4.4 L/ha	Apply at first appearance of disease and repeat at 7-10 day intervals.
Lettuce			
Beans			
Onions	Downy mildew	265-350ml per 100litres 3.2 - 4.4 L/ha	Apply when conditions favour infection and repeat at 7-10 day intervals.
Potatoes	Late blight, early blight	2.6 - 3.5 L/h	Apply every 7-10 days or more frequently under severe blight conditions. Increase the rate as plants increase in size.
Pumpkin, melons, cucumbers, other cucurbits	Downy mildew, anthracnose	265-350ml per 100litres 3.2 - 4.4 L/ha	Apply at 5-7 day intervals.
Tomatoes	Early blight, late blight, leaf mould, anthracnose	265-350ml per 100litres 4.4 L/ha	Apply at 7-10 day intervals or more frequently under blight conditions.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Etec Crop Solutions Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether incontract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

24 Hour Emergency Response
0800CHEM CALL
0800 243 622

Batch No:
DOM:

FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor if feeling unwell.

If spilt on skin, remove contaminated clothing and wash affected areas of skin. DO NOT SCRUB THE SKIN. If concentrate is splashed in eyes, flush with running water for at least 15 minutes. Take to hospital without delay. If inhaled, remove to fresh air. Take to hospital without delay.

SPILLAGE: Wear appropriate protective clothing. Absorb spills with sand, earth or other absorbent material and sweep up. Dispose of waste material at an approved landfill. Prevent liquid from entering sewers or bodies of water.

COMPATIBILITY: SuperManz is compatible with most commonly used agrichemicals. Do not mix with lime sulphur or Bordeaux mixture.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

WITHHOLDING PERIODS: Tomatoes, Cucurbits – 1 day. All other crops – 14 days.

PRODUCT INFORMATION: SuperManz is a flowable formulation of mancozeb. SuperManz is a protectant fungicide. Complete coverage is essential.

DIRECTIONS FOR USE:

It is an offence to use this product on animals

Mixing: Always shake the container before use. Pour the required amount into the partly filled spray tank with agitation running. Complete filling the tank. Maintain agitation until spraying is complete.



EL984



9 421022 422716



CMYK



Dieline

BACK Label