

**WARNING: KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# UniMaz<sup>TM</sup>

## 250 SL Herbicide

ACTIVE CONSTITUENT: 250 g/L IMAZAPYR  
present as the isopropylamine salt

**For the control of various annual and perennial weeds in  
non-crop situations as per the DIRECTIONS FOR USE table.**

**IMPORTANT: READ THE LABEL BEFORE USE**

GROUP B HERBICIDE



**AGRECOVERY**  
*Rural Recycling Programme*

*Approved pursuant to the HSNO Act 1996,  
Approval Code No. HSR100098*

**New Zealand Agent:**  
**ETEC Crop Solutions Limited**  
**P O Box 51584**  
**Pakuranga 2140 Auckland**  
**Phone 0800 100 325**  
**[www.etec.co.nz](http://www.etec.co.nz)**

NET  
CONTENTS:

**5**  
**Litres**

**Etec**<sup>®</sup>  
CROP SOLUTIONS

# PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and conditions of registration.

This product **MUST** be under the control of an approved handler when it is applied in a wide dispersive manner; or used by a commercial contractor.



**HSNO Classifications:**  
6.3A, 6.4A, 9.1A, 9.2B

## WARNING

May cause skin and eye irritation.

Very toxic to aquatic organisms.

Toxic to the soil environment

## PRECAUTIONS

Store in original container, tightly closed, away from foodstuffs and under lock and key.

Avoid contact with seeds, fertilisers and other pesticides.

When handling the concentrate wear full protective clothing such as overalls, gloves and face shield.

Avoid contact with skin, eyes and inhalation of spray mist.

Wash splashes of concentrate from skin or eyes immediately.

When spraying, wear overalls and gloves.

DO NOT eat, drink or smoke while using.

Wash hands and face before eating, drinking or smoking and after work.

Wash protective clothing after use.

Apply with well maintained and calibrated spray equipment.

Tracking records **MUST** be kept.

DO NOT carry on a passenger service vehicle.

Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.

Avoid contamination of any water supply with chemical or empty container.

UniMaz 250SL shall not be applied onto or into water.

**CONTAINER DISPOSAL:** Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, [www.agrecovery.co.nz](http://www.agrecovery.co.nz)). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

**PRODUCT DISPOSAL:** Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

**FIRST AID:** If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre **0800 POISON (0800 764 766)** or a doctor immediately. If skin, hair or eye contact occurs, remove contaminated clothing and flush skin, hair and eyes with running water for several minutes.

**SPILLAGE:** Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

## GENERAL INFORMATION

UniMaz 250 SL is a moderately residual herbicide for the control of annual and perennial grasses and broadleaved weeds on non cropland, non grazed areas. UniMaz 250 SL Herbicide is a member of the imidazolinone group of herbicides The active ingredient is rapidly absorbed through foliage and roots, and is translocated throughout plants.

Post emergence application is preferred. Foliage treatments will provide pre-emergence control as the active material is released to the soil following plant decay.

UniMaz 250 SL stops growth rapidly, although symptoms may take over 30 days to develop. Affected plants are discoloured and often growth is distorted. Residual activity may remain for up to 6 months after application.

## DIRECTIONS FOR USE

Apply the recommended rate for the weed species in sufficient water to avoid run off. For maximum activity, apply during periods of vigorous growth. Faster control of standing weeds will result if growing conditions are favourable at and after application.

Rainfall within two hours of the spray mixture drying on leaves may result in reduced weed control.

For the control of weeds around agricultural buildings and other farm non-crop non grazed situations, commercial, industrial, and public service areas, rights of way and waste land, away from desirable vegetation.	
WEEDS CONTROLLED AT 4 LITRES/HA	COMMENTS
Amaranthus, Annual Poa, Barnyard grass, Browntop, Buttercups, Carrot weed, Catsear, Chamomile, Chick weed, Cleavers, Clover, Couch, Cut-leaved geranium, Dandelion, Doves foot, Fathen, Fennel, Fescue, Field bindweed, Groundsel, Hedge mustard, Hemlock, Kikuyu, Mallows, Milkweed, Oxalis, Oxtongue, Oxeye daisy, Paspalum, Plantains, Prairy grass, Ryegrass, Scarlet pimpernel, Scotch Thistle, Sow thistle, Summer grass, Umbrella sedge, Vasey grass, Vetch, Willow weed, Wireweed, Witchgrass, Yarrow and Yorkshire fog	Apply 4 litres / ha in 50 – 200 litres water per hectare or sufficient to thoroughly wet the target weeds but avoiding run off.  For hand gun or knapsack application use 40ml / 100 litres of spray mix.
WEEDS CONTROLLED AT 6 LITRES/HA	COMMENTS
Black nightshade, Californian thistle, Canadian fleabane, Cockfoot, Docks, Hawksbeard, Indian doab, Mercer grass, Pampas grass, Soldiers button, Sorrel and Verbena	Apply 6 litres / ha in 50 – 200 litres water per hectare or sufficient to thoroughly wet the target weeds but avoiding run off.  For hand gun or knapsack application use 60ml / 100 litres of spray mix.

## DO NOT GRAZE OR CUT FOR STOCK FEED

**DO NOT graze:** Must not be applied to areas that maybe grazed by animals or cut for stock feed. Not to be used on or around food/ animal feed

## MIXING

Pour the required amount of UniMaZ 250 SL Herbicide into a spray container which already holds the total amount of water required.

Mix thoroughly. If UniMaZ 250 SL Herbicide is added during filling, foaming may occur. If excessive foaming is a problem, add a suitable antifoaming agent.

Consult your distributor for specific information.

UniMaZ 250 SL Herbicide may be applied in hard or soft water. The product is corrosive to mild steel.

Use ONLY stainless steel, fibreglass, plastic or plastic-lined containers for mixing, storage and application.

## COMPATIBILITY

For faster knockdown of weeds, certain other herbicides may be mixed with UniMaZ 250 SL Herbicide. Bipyridyl herbicides (eg. paraquat) are NOT compatible with UniMaZ 250 SL Herbicide. Mixtures of this type will result in poor weed control.

## SAFETY DATA SHEET

If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at [www.etec.co.nz](http://www.etec.co.nz)

## EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Etec Crop Solutions Ltd as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Unimaz is a trademark of United Phosphorus Ltd

Distributed by ETEC Crop Solutions

Approved pursuant to the HSNO Act 1996,  
Approval Code No. HSR100098.

UN 3082

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S.

(Contains 250 g/L imazapyr)

MARINE POLLUTANT

PACKING GROUP III, HAZCHEM 2X

In a transport emergency dial 111, Police or Fire Brigade



## 24 HOUR EMERGENCY RESPONSE

0800 CHEM CALL

0800 243 622

EL976

BATCH NO

DOM

