

CONTRA-INDICATIONS

A contra-indication is a condition that enables the client as unfit for this treatment or restricts you from performing this procedure. The condition may or may not be visible during your consultation so it is always important to discuss contra-indications with your client. Please also consult the following table of both contra-indications and cautions.

Contra - indications and special care

Positive reaction to a patch or sensitivity test (allergy). Patch test 24 hours before procedure	Do not perform treatment
Cysts (A cyst is a lump of the upper or lower eyelid caused by obstruction and inflammation of an oil gland of the eye lid - also known as a meibomian, tarsal or granuloma cyst)	Do not perform treatment
Hordeola/Style/Styes (a small collection of puss/abscess on the eyelid)	Do not perform treatment
Blepharitis (is an ocular condition characterized by chronic inflammation of the eyelid)	Do not perform treatment
Chemotherapy (cancer treatment)	Do not perform treatment
Skin Disease, skin trauma, cuts, abrasions, burns and swelling in the immediate area	Do not perform treatment
Weak eyelashes at therapist discretion on consultation (from over styling or product use)	Do not perform treatment
Skin Disorders in the general eye area (Dermatitis, Xanthelasma & Syringoma)	Do not perform treatment
Eye infections such as Iritis or conjunctivitis	Do not perform treatment
Eye inflammation such as Uveitis	Do not perform treatment
Recent operations around eyes, head or face or scar tissue in immediate areas	Do not perform treatment
Watery eyes	Do not perform treatment
Hypersensitive skin/eyes	Do not perform treatment
Keratitis (inflammation of the cornea of the eye)	Do not perform treatment
Alopecia (loss of hair)	Do not perform treatment
Trichotillomania (classified as an impulse control disorder and is the compulsive urge to pull out ones own hair)	Do not perform treatment
Bells Palsy or any condition that makes closing or opening eyes difficult	Do not perform treatment
Any disease/disorder that causes shaking, twitching or erratic movements	Do not perform treatment
Dry eye syndrome	Requires a GP referral
Glaucoma	Requires a GP referral
Post Chemotherapy	Requires a GP referral
Claustrophobia	Clients is required to not open eyes during treatment
Hayfever/Rhinitis	Eyes maybe watery - check client comfort
Contact lenses	Ensure removal before procedure
If using collagen eye pads during the lash lift treatment please ensure there is no previous history of reaction to collagen or the ingredients in the eye pad	Perform the treatment without the eye pads