


UN2992
 CARBANATE, PESTICIDE, LIQUID, TOXIC
 (contains furathiocarb 40%)
 PACKING GROUP II
 HAZCHEM 2XE



24 HOUR EMERGENCY RESPONSE
0800 CHEMCALL (0800 243 622)

TOXIC
KEEP OUT OF THE REACH OF CHILDREN
ECOTOXIC

Registered pursuant to the ACVM Act 1997 No. P8367 see www.foodsafety.govt.nz for registration conditions
 Approved pursuant to HSNO Act 1996 No. HSR000943 see www.epa.govt.nz for approval conditions

FURAKOTE EW

A seed treatment providing systemic protection of seedlings from attack by Argentine stem weevil on grasses and sweetcorn, nematodes on clovers, root knot nematode on carrots and springtails on brassicas

Contains 400g/litre furathiocarb in the form of an emulsion in water

NET CONTENTS: 200 Litres

FREE COLLECTION
 Drums 61-1000 litres
CONDITIONS APPLY
www.agrecovery.co.nz
 0800 AGRECOVERY

DISTRIBUTED BY
 Agrisource (2000) Ltd
 45 Kitchener Road
 Pukekohe 2120
 Ph: 09 239 0275
www.agrisource.co.nz

BATCH NO:

DATE OF
 MANUFACTURE:

READ LABEL AND SAFETY DATA SHEET COMPLETELY BEFORE USE

HSNO CLASSIFICATIONS: 6.1B, 6.1C, 6.3B, 6.4A, 6.5B, 6.9A, 9.1A, 9.2B, 9.3A, 9.4A

APPROVED HANDLER

This product must be under the personal control of an approved handler or secured.

TRACKING

Location and movement of this product must be recorded at each stage of its lifecycle.

TOXIC

May be fatal if swallowed, inhaled or absorbed through the skin. May cause target organ damage from repeated oral exposure at high doses.

HARMFUL

May cause skin and eye irritation. May cause sensitisation from prolonged skin contact. Avoid skin or eye contact.

ECOTOXIC

Very toxic to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container.
Toxic to the soil environment. Very toxic to terrestrial vertebrates, Very Toxic to terrestrial invertebrates.

HANDLING PRECAUTIONS

Personal Protection: When mixing or applying, avoid contact with skin and eyes. Wear full protective clothing, gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work.

Storage: Store in original container tightly closed and in a locked, dry, cool, well ventilated area away from fertilisers, feed, seeds and foodstuffs. Keep out of reach of children. **Avoid temperatures below 0°C and above 35°C.**

Under HSNO regulations storage of certain quantities of this product may require bunding & signage. See the PRODUCT SAFETY CARD for further information.

Shelf Life: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this. **After prolonged storage >12 months, it is advisable to stir or shake the drum to ensure there is no separation of product, before opening and using the contents.**

Spillage: Move non-essential people from the area. Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Product Disposal: Dispose of unwanted product is by used by another operator strictly in accordance with label directions.

Container Disposal: Triple rinse drum and add rinsate to mixing tank. Contact the Agrecovery rural recycling programme for free drum collection on 0800 AGRECOVERY (0800 247 326). Do not use container for any other purpose.

Safety Data Sheet: Additional information is provided in the SDS available from your supplier.

FIRST AID

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 Poison (0800764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If concentrate is splashed in the eyes, flush with running water immediately.

EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade.

PRODUCT INFORMATION

FURAKOTE EW is a systemic insecticide for seed treatment only. It is absorbed via the emerging young roots and transported via the sap stream upwards in the seedling plant. Approximately 4 weeks protection should be achieved against soil inhabiting pests.

It is an offence to use this product on animals.

DIRECTIONS FOR USE

EQUIPMENT

Apply **FURAKOTE EW** using suitable, clean batch treatment equipment. Clean equipment thoroughly with water and detergent after use before residues can dry.

COMPATIBILITY

FURAKOTE EW may be mixed with other seed treatment compounds.

APPLICATION METHOD

Apply to clean seed only, use **FURAKOTE EW** undiluted, apply to seed allowing minimal mixing time.

Note: on waxy seeds such as Maize and Sweetcorn a suitable polymer binder will improve adherence. With these seeds it is recommended to add a high-grade talcum powder at 1kg/80,000 seeds to improve seed flow characteristics and drying, allow 3 days curing before planting.

Treated seed must be permitted to 'dry off' - don't pack into plastic lined bags. When seed is artificially dried, the temperature of the seed must not exceed 35°C.

TREATED SEED

Must be dyed a distinctive colour and containers of seed must be labeled with the following statement:

FURAKOTE EW treated seed must NOT be used for human or animal consumption.

FURAKOTE EW treated seed maybe harmful to seed eating birds, it must be sown properly and completely covered by soil.

FURAKOTE EW treated seed is NOT to be left in areas accessible to birds.

WITHHOLDING PERIODS

Stock must not graze grasses, clovers or brassicas within 42 days of sowing, or maize and sweetcorn within 60 days of sowing.

CROP	PEST	RATE	REMARKS
Brassicas	Springtails	12.5ml/kg of seed	Refer Application method
Carrots	Root Knot Nematodes	100ml/kg of seed	Refer Application method
Clovers	Nematodes	50ml/kg of seed	Refer Application method
Grasses	Argentine Stem Weevil larvae	25ml/kg of seed	Refer Application method
Grain & Silage Maize	Argentine Stem Weevil	750 ml/ 80,000seeds	Refer Application method
Sweetcorn	Argentine Stem Weevil	30-55ml/kg of seed	Use 30ml rate on seed counts of 3,000/kg, use 55ml rate where maximum count is 6,000 seeds/kg, use appropriate rates in between

It is an offence for the users of this product to cause residues exceeding the relevant MRL in the NZ (Maximum Residues Limits of Agricultural Compounds) Food Standards.

CONDITIONS OF SUPPLY

All goods supplied by us are produced under rigid quality control procedures which assure the purchaser that at the time of sale they are physically suitable for use in the manner stated. However, to the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agri Solutions Ltd as registrant, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

REGISTRANT: Agri Solutions Ltd, Level 1, 26 Crummer Road, Grey Lynn, Auckland 1021