



**AGRISOURCE**  
PRODUCTS FOR AGRICULTURE

**HARMFUL**  
**KEEP OUT OF THE**  
**REACH OF CHILDREN**

**ECOTOXIC**

# Delete

**Selective post-emergence herbicide for use on cereals,  
clover and grass seed crops, dwarf beans, lucerne, onions,  
pasture, peas, potatoes, and turf.**

*Contains 480g/litre bentazone as the sodium salt in the form of a soluble concentrate*

GROUP	<b>C3</b>	HERBICIDE
-------	-----------	-----------



**NET CONTENTS**  
**10 Litres**

# READ LABEL COMPLETELY BEFORE USE

## DIRECTIONS FOR USE

NOTE: Wetting agents should not be added to Delete unless where specified.

Allow at least 8 hours drying before rainfall or irrigation. Delete is not recommended for Low Volume CDA application.

**HSNO CLASSIFICATIONS: 6.1D, 6.4A, 6.5B, 9.1C, 9.3C**

### APPROVED HANDLER

This product must be under the control of an approved handler during use.



### HARMFUL

This substance may be harmful if swallowed, inhaled or absorbed through the skin. May cause eye irritation. Avoid eye contact. May cause sensitisation from prolonged skin contact.

### ECOTOXIC

Harmful to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to terrestrial vertebrates.

## HANDLING PRECAUTIONS

**Personal Protection:** When mixing or applying, avoid contact with skin and eyes. Wear full protective clothing, gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work.

**Equipment:** Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409). Flush spray equipment thoroughly with water after use.

**Storage:** Store in original container tightly closed and in a locked, dry, cool, well ventilated area away from fertilisers, feed, seeds and foodstuffs.

Keep out of reach of children. Store in accordance with NZ 8409 Management of Agrichemicals.

**Shelf Life:** When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

**Spillage:** Move non-essential people from the area. Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

**Product Disposal:** Dispose of unwanted product is by used by another operator strictly in accordance with label directions.

**Container Disposal:** Triple rinse container and add rinsate to spray tank. Dispose of cleaned container at your local AGRECOVERY container collection site.

**Safety Data Sheet:** Additional information is provided in the SDS available from your supplier.

## FIRST AID

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If concentrate is splashed in the eyes, flush with running water immediately.

## EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade

## GENERAL DIRECTIONS

**Delete** is a selective contact post-emergence herbicide recommended for the control of broad-leaved weeds in many crops.

**Delete** must be applied in sufficient water to ensure that all parts of the target weeds are thoroughly covered. Immature weeds (2-4 leaf) are more readily controlled than more mature plants.

**It is an offence to use this product on animals.**

## IMPORTANT - RESISTANCE MANAGEMENT

**Delete** contains the **GROUP 25 HERBICIDE** benazotone. Resistance to pesticides can develop from excessive use. To minimise this risk, avoid using this herbicide exclusively all season, avoid unnecessary spraying and endeavour to alternate with herbicides from other groups within the crop and in subsequent seasons for a sprayed paddock. Use strictly in accordance with label instructions. Maintain good cultural practices.

## WEEDS CONTROLLED

**Delete** is especially effective against chamomiles, stinking mayweed, cleavers, storksbill and Onehunga weed. Other weeds controlled include redroot (Prince of Wales feather), chickweed, black nightshade, spurry (yarr), shepherds purse, thorn apple, and wild turnip.

## WHEN TO APPLY

**Delete** is not active in the soil, so weeds must have germinated before spraying (post emergence).

## MIXING

Half fill spray tank with water, with agitation running, add the required amount of **Delete**, top up with water and agitate thoroughly before use.

CROP	TIMING	RATE	REMARKS
Cereals	After 4 true leaves	2L/ha in 170-500L of water.	Apply in combination with sodium salt formulations of MCPB, dichlorprop or mecoprop, applied at label recommended rates. Delete + MCPB may be applied to crops under sown with clover.
Clover Seed Crops	In late winter early spring, when seeds are small	2-3L/ha in a minimum of 500L of water.	Use high water rates and large droplets to ensure penetration into growth crown of weeds. Add a non-ionic spreader/sticker Paraquat, applied at least 14 days before Delete treatment will improve results, provided weeds are actively growing at the time of Delete treatment. Spraying during warm weather gives best results.
Dwarf Beans	When crop has 1-2 true leaves and weeds are at cotyledon to 4 leaf stage	2-3L/ha in a minimum of 500L of water.	Some leaf scorch may occur, but yield is unlikely to be affected. Do not apply to crops before 1 true leaf stage and avoid conditions above 21C. Use higher rates on more mature weeds. Add a non-ionic spreader-sticker to enhance activity on Thorn apple and hairy nightshade.
Grass Seed Crops	When weeds are at a susceptible stage	2-3L/ha in a minimum of 500L of water.	Ensure thorough coverage of weeds. Add a non-ionic spreader/sticker. Spraying during warm weather gives best results.
Lucerne	During Autumn	1.5L plus 2L of 2, 4-DB (40% a) per ha. Or 4 L/ha alone. Apply in 300-500L of water.	For control of nodding thistle and storksbill.
New Pasture	When clover has 1-2 true leaves and weeds are at a 4-6 leaf stage	3L/ha in 300-500L of water.	Delete may be applied as a tank mix with MCPB.
Onions	After onions have 3 leaves	2-3L/ha in 330-550L of water.	Apply when weeds are still seedlings and onions are waxy and healthy. Do not add surfactants. Do not apply when soil is water logged. Do not apply when onions are damaged by hail, frost, heavy rain, wind etc. Do not apply other herbicides for 10 days before or after Delete application.
Peas	When crop has 1-2 true leaves and weeds are at 4 leaf stage	2L/ha plus 2.5L/ha MCPB Apply in 300-500L of water.	Suitable for use on under sown pea crops.
Potatoes	When cleavers are visible and prior to crop emergence	2-3L/ha in 300-500L of water	
Turf	October to early November	3L/ha in 300-500 L of water. Or 30ml/100 square metres in 3-5L.	A second Autumn spray will reduce the problem for next season. In specialised equipment, e.g. "Lawnboy".

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

## CONDITIONS OF SUPPLY

All goods supplied by us are produced under rigid quality control procedures which assure the purchaser that at the time of sale they are physically suitable for use in the manner stated. However, to the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agri Solutions Ltd as registrant, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered pursuant to the ACVM Act 1997 No. P8601  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Approved pursuant to HSNO Act 1996 No. HSR000426  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

## REGISTRANT

Agri Solutions Ltd  
Level 1, 26 Crummer Road  
Grey Lynn, Auckland 1021

## DISTRIBUTOR

Agrisource (2000) Ltd  
45 Kitchener Road  
Pukekohe 2120  
09 237 0442 · [www.agrisource.co.nz](http://www.agrisource.co.nz)

BATCH NO:

DATE OF  
MANUFACTURE:

# 24 HOUR EMERGENCY RESPONSE 0800 CHEMCALL (0800 243 622)