

TexaClear® Kids Allergy Relief

Drug Facts

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each 30 mL (2 TBSP))	Purpose
Chlophedianol HCl 12.5 mg	Cough Suppressant
Pyrilamine Maleate 25 mg	Antihistamine
Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies: • cough due to minor throat and bronchial irritation • runny nose • sneezing • itchy, watery eyes • itching of the nose or throat	
Warnings Do not exceed recommended dosage.	
Ask a doctor before use if you have <ul style="list-style-type: none"> ▪ a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema ▪ glaucoma ▪ difficulty in urination due to enlargement of the prostate gland ▪ cough that occurs with too much phlegm (mucus) ▪ a breathing problem such as emphysema or chronic bronchitis 	
Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers	
When using this product <ul style="list-style-type: none"> ▪ do not use more than directed ▪ marked drowsiness may occur ▪ alcohol, sedatives, and tranquilizers may increase drowsiness ▪ avoid alcoholic drinks ▪ be careful when driving a motor vehicle or operating machinery ▪ excitability may occur, especially in children 	
Stop use and ask a doctor if <ul style="list-style-type: none"> ▪ nervousness, dizziness, or sleeplessness occurs ▪ symptoms do not improve within 7 days, tend to recur, or are accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition. <ul style="list-style-type: none"> ▪ new symptoms occur 	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none"> ▪ do not take more than 4 doses in any 24-hour period. ▪ do not exceed recommended dosage. ▪ use enclosed dose cup ▪ keep dosage cup with product ▪ TBSP= tablespoon, mL= milliliter 	
adults and children 12 years of age and over:	60 mL (4 TBSP) every 6 to 8 hours
children 6 to under 12 years of age:	30 mL (2 TBSP) every 6 to 8 hours
children under 6 years of age:	do not use.
Other information <ul style="list-style-type: none"> ▪ each 30 mL (2 TBSP) contains: Sodium 23 mg ▪ tamper evident: Do not use if foil seal under cap is broken or missing. ▪ read all product information before using ▪ store at room temperature 68-86°F (20-30°C) ▪ avoid excessive heat and humidity 	
Inactive ingredients bitter mask, citric acid anhydrous, glycerin, polyethylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose	
Questions or comments? Call 1-888-535-0305 9 a.m. - 5 p.m. CST.	



Distributed by:
GM Pharmaceuticals, Inc.
Arlington, TX. 76015