

# TexaClear® Kids Daytime Cough & Cold Relief Drug Facts

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

<b>Drug Facts</b>	
<b>Active ingredients (in each 30 mL (2 TBSP))</b>	<b>Purpose</b>
Dextromethorphan HBr 10 mg.....	Cough Suppressant
Guaifenesin 200 mg .....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>▪ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.</li> <li>▪ temporarily relieves: ▪ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ▪ the intensity of coughing ▪ nasal congestion due to a cold ▪ stuffy nose</li> </ul>	
<b>Warnings</b>	
Do not exceed recommended dosage.	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>▪ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>▪ in children under 6 years of age</li> </ul>	
<b>Ask a doctor before use if you have</b>	
<ul style="list-style-type: none"> <li>▪ heart disease, high blood pressure, thyroid disease, or diabetes</li> <li>▪ cough that occurs with too much phlegm (mucus)</li> <li>▪ persistent or chronic cough such as occurs with asthma, smoking, chronic bronchitis, or emphysema</li> <li>▪ trouble urinating due to an enlarged prostate gland</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are</b> ▪ taking any other oral nasal decongestant or stimulant	
<b>When using this product ▪ do not use more than directed</b>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>▪ symptoms do not improve within 7 days or are accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.</li> <li>▪ nervousness, dizziness, or sleeplessness occurs</li> </ul>	
<b>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>▪ do not take more than 6 doses in any 24-hour period.</li> <li>▪ do not exceed recommended dosage.</li> <li>▪ use enclosed dose cup</li> <li>▪ keep dosage cup with product</li> <li>▪ TBSP= tablespoon, mL= milliliter</li> </ul>	
adults and children 12 years of age and over:	60 mL (4 TBSP) every 4 hours
children 6 to under 12 years of age:	30 mL (2 TBSP) every 4 hours
children under 6 years of age:	do not use.
<b>Other information</b>	
<ul style="list-style-type: none"> <li>▪ each 30 mL (2 TBSP) contains: Sodium 20 mg</li> <li>▪ tamper evident: Do not use if foil seal under cap is broken or missing.</li> <li>▪ read all product information before using</li> <li>▪ store at room temperature 68-86°F (20-30°C)</li> <li>▪ avoid excessive heat and humidity</li> </ul>	
<b>Inactive ingredients</b>	
citric acid anhydrous, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose	
<b>Questions or comments?</b>	
Call 1-888-535-0305 9 a.m. - 5 p.m. CST.	



Distributed by:  
GM Pharmaceuticals, Inc.  
Arlington, TX. 76015