

Optoelectronics (광전자공학)

Lecture 3. Optical properties of solids

Young Min Song

Associate Professor

School of Electrical Engineering and Computer Science


Gwangju Institute of Science and Technology

<http://www.gist-foel.net>

ymsong@gist.ac.kr, ymsong81@gmail.com


A207, ☎2655

Optical coefficients




- Reflectivity = reflected / incident power
- Transmissivity = transmitted / incident power
- $T + R = 1$ if medium is transparent

- Luminescence comes out at lower frequency than absorption due to internal relaxation.
- The energy shift between absorption and luminescence is called the Stokes shift.



Optical coefficients

propagation through
the medium





Optical coefficients

Complex refractive index


Complex refractive index

Insulators/semiconductors




- Infrared absorption due to phonons
- UV/visible absorption due to bound electrons
- Position of fundamental absorption edge depends on the bandgap
- Cr^{3+} ions doped into sapphire absorb in the blue and yellow/green spectral regions, hence red colour

Metals/Organic materials



- Free electrons in the metal absorb
→ High reflectivity up to 'plasma frequency' in the UV




- Strong absorption in UV/visible spectral region due to electronic transitions
- Stokes-shifted emission across the visible spectral region

Dipole oscillators



Classical model of an atom. Electrons are bound to the nucleus by springs which determine the natural frequencies.




Dipole oscillators

Dipole oscillators

Dipole oscillators


Dipole oscillators



Lorentz oscillator with

- $\omega_0 = 10^{14}$ rad/s
- $\gamma = 5 \times 10^{12}$ s⁻¹
- $\epsilon_{st} = 12.1$
- $\epsilon_\infty = 10$

Atomic absorption line




Lorentzian lineshape

$$\alpha(\omega) = \alpha_0 \frac{\gamma^2}{4\Delta\omega^2 + \gamma^2}$$


$$n(\omega) = n_0 - \Delta n \frac{4\gamma\Delta\omega}{4\Delta\omega^2 + \gamma^2}$$

$$\Delta\omega = \omega - \omega_0$$

Multiple resonances




SiO₂ glass



Kramers-Kronig relationships

$$n(\omega) - 1 = \frac{2}{\pi} \text{P} \int_0^{\infty} \frac{\omega' \kappa(\omega')}{\omega'^2 - \omega^2} d\omega'$$

$$\kappa(\omega) = -\frac{2}{\pi\omega} \text{P} \int_0^{\infty} \frac{\omega'^2 [n(\omega') - 1]}{\omega'^2 - \omega^2} d\omega'$$




UV transmission of glass



Table 1.4 Composition, refractive index and ultraviolet transmission of some common glasses. The letters after the names give the abbreviations used to identify the glass type. The composition figures are the percentage by mass. The refractive index is measured at 546.1 nm, and the transmission is for a 1 cm plate at 310 nm. (Data from Driscoll & (1978), and Lide (1996).)

Name	SiO ₂	B ₂ O ₃	Al ₂ O ₃	Na ₂ O	K ₂ O	CaO	BaO	PbO	P ₂ O ₅	<i>n</i>	<i>T</i>
Fused silica	100									1.460	0.91
Crown (K)	74			9	11	6				1.513	0.4
Borosilicate crown (BK)	70	10		8	8	1	3			1.519	0.35
Phosphate crown (PK)		3	10		12	5			70	1.527	0.46
Light flint (LF)	53			5	8			34		1.585	0.008
Flint (F)	47			2	7			44		1.607	–
Dense flint (SF)	33				5			62		1.746	–


Dispersion



Double refraction (Birefringence)




Polarizing beam splitters



Additional : Plasma reflectivity

- Plasma : a neutral gas of charged particles (e.g., metals and doped semiconductors)
- The free electrons experiences no restoring forces when they interact with EM waves. \leftrightarrow Bound electrons


Additional : Plasma reflectivity




Additional : Free carrier conductivity

Additional : Free carrier conductivity

Additional : Metals



Additional : Doped semiconductors



Question or Comment?