

Checklist for prescribing opioids for chronic pain

For primary care providers treating adults (18+) with chronic pain ≥ 3 months, excluding cancer, palliative, and end-of-life care

CHECKLIST

When **CONSIDERING** long-term opioid therapy

- Set realistic goals for pain and function based on diagnosis (eg, walk around the block).
- Check that non-opioid therapies tried and optimized.
- Discuss benefits and risks (eg, addiction, overdose) with patient.
- Evaluate risk of harm or misuse.
 - Discuss risk factors with patient.
 - Check prescription drug monitoring program (PDMP) data.
 - Check urine drug screen.
- Set criteria for stopping or continuing opioids.
- Assess baseline pain and function (eg, PEG scale).
- Schedule initial reassessment within 1–4 weeks.
- Prescribe short-acting opioids using lowest dosage on product labeling; match duration to scheduled reassessment.

If **RENEWING** without patient visit

- Check that return visit is scheduled ≤ 3 months from last visit.

When **REASSESSING** at return visit

Continue opioids only after confirming clinically meaningful improvements in pain and function without significant risks or harm.

- Assess pain and function (eg, PEG); compare results to baseline.
- Evaluate risk of harm or misuse:
 - Observe patient for signs of over-sedation or overdose risk.
 - If yes: Taper dose.
 - Check PDMP.
 - Check for opioid use disorder if indicated (eg, difficulty controlling use).
 - If yes: Refer for treatment.
- Check that non-opioid therapies optimized.
- Determine whether to continue, adjust, taper, or stop opioids.
- Calculate opioid dosage morphine milligram equivalent (MME).
 - If ≥ 50 MME/day total (≥ 50 mg hydrocodone; ≥ 33 mg oxycodone), increase frequency of follow-up; consider offering naloxone.
 - Avoid ≥ 90 MME/day total (≥ 90 mg hydrocodone; ≥ 60 mg oxycodone), or carefully justify; consider specialist referral.
- Schedule reassessment at regular intervals (≤ 3 months).

REFERENCE

EVIDENCE ABOUT OPIOID THERAPY

- Benefits of long-term opioid therapy for chronic pain not well supported by evidence.
- Short-term benefits small to moderate for pain; inconsistent for function.
- Insufficient evidence for long-term benefits in low back pain, headache, and fibromyalgia.

NON-OPIOID THERAPIES

Use alone or combined with opioids, as indicated:

- Non-opioid medications (eg, NSAIDs, TCAs, SNRIs, anti-convulsants).
- Physical treatments (eg, exercise therapy, weight loss).
- Behavioral treatment (eg, CBT).
- Procedures (eg, intra-articular corticosteroids).

EVALUATING RISK OF HARM OR MISUSE

Known risk factors include:

- Illegal drug use; prescription drug use for nonmedical reasons.
- History of substance use disorder or overdose.
- Mental health conditions (eg, depression, anxiety).
- Sleep-disordered breathing.
- Concurrent benzodiazepine use.

Urine drug testing: Check to confirm presence of prescribed substances and for undisclosed prescription drug or illicit substance use.

Prescription drug monitoring program (PDMP):

Check for opioids or benzodiazepines from other sources.

ASSESSING PAIN & FUNCTION USING PEG SCALE

PEG score = average 3 individual question scores (30% improvement from baseline is clinically meaningful)

Q1: What number from 0–10 best describes your **pain** in the past week?

0 = “no pain”, 10 = “worst you can imagine”

Q2: What number from 0–10 describes how, during the past week, pain has interfered with your **enjoyment of life**?

0 = “not at all”, 10 = “complete interference”

Q3: What number from 0–10 describes how, during the past week, pain has interfered with your **general activity**?

0 = “not at all”, 10 = “complete interference”



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

TO LEARN MORE

www.cdc.gov/drugoverdose/prescribing/guideline.html